

# DUKE MATH MEET 2016

## TEAM SOLUTIONS

1. It is easy to show that  $\boxed{8}$  is possible. First each tile has area 4 and total area is 36 so we can have at most 9 tiles. We will show that 9 is not possible. Color the  $6 \times 6$  grid in a checkboard pattern. Then there are an 18 white squares. Each tile will cover either 1 or 3 white squares. Hence 9 tiles will cover an odd number of white squares. This isn't possible so the maximum is 8.
2.  $\angle BEC = 90^\circ$  and  $\angle CDB = 90^\circ$ . So  $BECD$  is a cyclic quadrilateral. Let  $F$  be the intersection of  $BD$  and  $CE$ . Then  $\triangle DEF$  is similar to  $BCF$ . Hence  $\frac{DE}{BC} = \frac{FD}{BF}$  but triangle  $BDF$  is a 30-60-90 right triangle so  $\frac{FD}{BF} = \frac{\sqrt{3}}{2}$ .
3. So we have  $2f(x) + f(1-x) = x^2$  and  $2f(1-x) + f(x) = (1-x)^2$  (we substitute  $1-x$  for  $x$ ). We can solve this as a system of linear equations. If we multiply the first equation by 2 and then subtract the second, we see that  $3f(x) = 2x^2 - (1-x)^2 = x^2 - 2x - 1$ . Hence the sum of the coefficients is  $\frac{1}{9}(1 + 4 + 1) = \frac{2}{3}$ .
4. We want to find the minimum integer  $k$  of the form  $15m^2 - a^2$  where  $15m^2 < a^2 - 1$ . Checking  $k = 0, \dots, 5$ , all will not satisfy through modulo 3, 2, and 5.  $15m^2 - 6 = a^2$ . So  $a = 3b$  so we have  $5m^2 - 2 = 3b^2$ . Modulo 5, we see that  $b \equiv 1, 4 \pmod{5}$ . Trying possibilities, we see that  $b = 9, m = 7$  works. So the answer is  $\boxed{6}$ .
5. Expanding  $(\sqrt{5} + 2)^{2016} + (\sqrt{5} - 2)^{2016}$ , we see that it is equal to an integer. In addition  $\sqrt{5} - 2 < 1$  so any power of it is less than 1. So  $\lfloor (\sqrt{5} + 2)^{2016} \rfloor = (\sqrt{5} + 2)^{2016} + (\sqrt{5} - 2)^{2016} - 1$ . Since we only need the last two digits, we can consider the expression mod 100.  $(\sqrt{5} + 2)^{2016} + (\sqrt{5} - 2)^{2016} = 2(5)^{1008} + 2\binom{2016}{2}(5)^{1007}2^2 + \dots + \binom{2016}{2}2(5)2^{2014} + 2(2)^{2016}$ . Note that most terms are divisible by 100 so we can ignore them. So we have  $2(5)^{1008} + 2^{2017}$ .  $2^{2017}$  repeats every  $\phi(25) = 20 \pmod{25}$  so we have  $2^{2017} \equiv 2^{17} \equiv 2^{-3} \equiv -3 \pmod{25}$ . So  $2^{2017} \equiv 72 \pmod{100}$ .  $2(5)^{1008} + 2^{2017} \equiv 50 + 72 \equiv 22 \pmod{100}$ . Hence  $22 - 1 = \boxed{21}$ .
6. Suppose  $f(2^a 3^b)$  is the maximum over the given range with  $2^a 3^b$  smallest. Since  $f(2^{a-5} 3^{b+3}) = f(2^a 3^b)$  but  $2^{a-5} 3^{b+3} = \frac{27}{32} 2^a 3^b$ . Then we see that  $a < 5$  otherwise we contradict our minimality of  $2^a 3^b$ . For a fixed  $a$ , we just want to pick the largest  $b$  such that  $2^a 3^b \leq 10000$ . For powers of 3, we have 1, 3, 9, 27, 243, 729, 2187, 6561. So when  $a = 0$ , we get  $b = 8$ .  $a = 1 \implies b = 7$ ,  $a = 2 \implies b = 7$ ,  $a = 3 \implies b = 6$ ,  $a = 4 \implies b = 5$ . Calculating  $3a + 5b$  we see that the maximum is when  $a = 2, b = 7$  so  $3(2) + 7(5) = \boxed{41}$ .
7. Considering only  $4n + 3$  primes. Suppose  $x$  is any odd number and  $p$  a  $4n + 3$  prime. Then  $x, px, p^2x, \dots, p^{2n-1}x$  contains the same number of  $4n + 1$  numbers as  $4n + 3$ . So we see that every power of a  $4n + 3$  prime must be even. In addition, we can see that

when all the  $4n + 3$  primes are raised to an even power that is has precisely 1 more  $4n + 1$  divisor than  $4n + 3$ . Now consider the  $4n + 1$  prime factors. Let  $x$  be the product of all  $4n + 1$  prime factors. Since multiplying by  $4n + 1$  doesn't change if the number is  $4n + 1$  or  $4n + 3$ , we see that multiplying by  $x$  to a product of  $4n + 3$  prime powers just multiplies the difference in  $4n + 1$  and  $4n + 3$  powers by the number of divisors of  $x$ . So we just need  $x$  to have 6 divisors ( $5^5$  works) and the  $4n + 3$  powers to be even. So a possible answer is  $\boxed{5^5 3^{27^2}}$ .

8. Consider the graph  $y = x^{3/2}$ .  $\lfloor i^{3/2} \rfloor$  is the number of lattice points below the graph at  $x = i$  (includes the point on the graph if  $i^{3/2}$  is an integer. But  $y = x^{2/3}$  is the inverse of  $y = x^{3/2}$ . So  $\lfloor i^{2/3} \rfloor$  is the number of lattice points to the left of the graph of  $y = x^{3/2}$  at  $y = i$ . Hence overall since the bounds match up, this is just the area of a rectangle with side lengths 100 and 1000 plus the number of lattice points on the graph which is just 10. Hence our answer is  $100(1000) + 10 = \boxed{100010}$ .

9. We find the probability that  $A \subseteq B$ . The probability that an element is in  $A$  but not in  $B$  is  $\frac{1}{4}$ . So the probability that  $A \subseteq B$  is  $\frac{3^{10}}{4^{10}}$ . But we are overcounting the probability that  $A = B$ . The probability that an element is in  $A$  and in  $B$  is  $\frac{1}{2}$ . So probability that  $A = B$  is  $\frac{1}{2^{10}}$ . So overall probability is  $\frac{2(3^{10}) - 2^{10}}{4^{10}}$ .

10. The answer is 21. Suppose there are more than 21 teams. Let the teams be  $A_1, \dots, A_k$ . Then  $|A_i| = 5$  and  $|A_i \cap A_j| = 1$ . Consider the intersection of  $A_1$  with  $A_2, \dots, A_k$ . Then some element of  $A_1$  must appear at least 5 times by pigeonhole principle. Hence we have at least six teams sharing a person  $a$ . Call these teams  $B_1, B_2, B_3, B_4, B_5, B_6$ . Let  $b$  be a person on  $B_1$  and  $b'$  in  $B_2$ . This team must intersect  $B_3, B_4, B_5, B_6$  but this is not possible since the only element these sets share is  $a$ . So we can have at most 21 teams.

On the other hand, if there are less than 21 teams, some person  $a$  can be on at most 4 teams. Suppose these are  $B_1, B_2, B_3, B_4$ . Take some person  $b$  from  $B_1$  and  $b'$  from  $B_2$ . Then this team must share a teammate with  $B_3$  and  $B_4$ . The last teammate  $c$  cannot come from  $B_1, B_2, B_3, B_4$  otherwise two teams have the same pair of people. But then  $a$  and  $c$  are never on a team.

So the answer is  $\boxed{21}$ .