


PUMaC 2008-9



Number Theory - A

1. What is the remainder, in base 10, when $24_7 + 364_7 + 43_7 + 12_7 + 3_7 + 1_7$ is divided by 6?

(ANS: 3 CB: GBH?, ACH, KB)

2. How many zeros are there at the end of $792!$ when written in base 10?

(ANS: 196 CB: ACH)

3. Find all integral solutions to $x^y - y^x = 1$.

(ANS: (2, 1) and (3, 2) must be given; $(x \neq 0, 0)$ may also be. If both x and y are greater than 3 and not equal, we have $|x^y - y^x| > 2$ for analytic reasons. Thus one of x and y are 1 or 2, so you can check that the two given are the only solutions.

In some versions of the test, the word “positive” was included; therefore the third solution (with $y = 0$) was optional. Credit was given to any contestants who answered either version correctly, that is, who gave either all three answers or just the first two. CB: AL, IAF)

4. Find the largest integer n , where 2009^n divides $2008^{2009^{2010}} + 2010^{2009^{2008}}$.

(ANS: Using the binomial theorem:

$$A = (2009 - 1)^{2009^{2010}} = -1 + 2009 * (2009^{2010}) + R_1$$

$$B = (2009 + 1)^{2009^{2008}} = 1 + 2009 * (2009^{2008}) + R_2$$


R_1 is divisible by 2009^{2010} and R_2 is divisible by 2009^{2010} . $A + B \equiv 2009^{2009} \pmod{2009^{2010}}$, so $n = 2009$. CB: AL)

5. How many integers n are there such that $0 \leq n \leq 720$ and $n^2 \equiv 1 \pmod{720}$?

(ANS: 16. $720 = 16 * 9 * 5$. There are two solutions mod 5 and mod 9, and there are 4 mod 16, and so, by the chinese remainder theorem, there are 16 solutions mod 720. But the numbers $1, 2, \dots, 720$ form a complete residue system mod 720, so there are 16 solutions in all. CB: IAF)

6. $f(n)$ is the sum of all integers less than n and relatively prime to n . Find all integers n such that there exist integers k and l such that $f(n^k) = n^l$.

PUMaC 2008-9



Number Theory - A

(ANS: 2, 3, 4, 6. Let $\phi(n)$ denote the number of positive integers less than n that are relatively prime to n . It is clear that $2f(n) = n\phi(n)$, and so we need either $\phi(n) = 1$ or $\phi(n) = 2$, and one can check that 2, 3, 4, and 6 are the only solutions to this. Moreover, for those values of n , we have $f(n) = n$ (except 2, where we have $f(2) = 1$) **CB: AL, IAF**)

7. In this problem, we consider only polynomials with integer coefficients. Call two polynomials p and q *really close* if $p(2k+1) \equiv q(2k+1) \pmod{2^{10}}$ for all $k \in \mathbb{Z}^+$. Call a polynomial p *partial credit* if no polynomial of lesser degree is *really close* to it. What is the maximum possible degree of partial credit?

(ANS: 5.


Suppose we have a polynomial $p(n)$ that has the maximum possible degree of partial credit. Consider $q(x) = (x-1)(x-3)(x-5)(x-7)(x-9)(x-11)$. Observe that for all odd integers x , $q(x)$ is the product of six consecutive even integers, and thus is divisible by $2 \cdot 4 \cdot 6 \cdot 8 \cdot 10 \cdot 12$. Note that the power of two dividing $2 \cdot 4 \cdot 6 \cdot 8 \cdot 10 \cdot 12$ is 2^{10} . Thus we conclude that $q(2k+1) \equiv 0 \pmod{2^{10}}$. Now if $p(n)$ has degree six or larger, let its leading term be an^ℓ . Then $p(n)$ and $p(n) - an^{\ell-6}q(n)$ are really close, but the latter has lesser degree. Thus the maximum possible degree is at most five.

Consider $p(x) = (x-1)(x-3)(x-5)(x-7)(x-9)$. This is a polynomial of degree five. I claim that this polynomial is partial credit. Suppose for sake of contradiction that there exists a polynomial $q(x)$ of degree four or less such that $p(2k+1) \equiv q(2k+1) \pmod{2^{10}}$ for all k . We know $q(1)$ is divisible by 1024. We may add a multiple of 1024 to $q(x)$ so that $q(1) = 0$ and q is still really close to p . Thus $q(x) = (x-1)r(x)$. Now consider $q(3)$. It is divisible by 1024, so $r(3)$ is divisible by 512. Now adding a multiple of 512 to $r(x)$ does not change the residue class of $q(x)$ modulo 1024 for x odd. Thus we may assume that $r(3) = 0$. Hence we get $q(x) = (x-1)(x-3)s(x)$. Similarly, we find $s(5)$ is divisible by 128, and adding multiples of 128 to $s(x)$ does not change the residue class of $q(x)$ modulo 1024 for odd values of x . Hence WLOG $q(x) = (x-1)(x-3)(x-5)h(x)$. Iterating this argument, we find $q(x) = (x-1)(x-3)(x-5)(x-7)(x-9)k(x)$ for some polynomial $k(x)$. Since q has degree less than five, we conclude that k is the zero polynomial. Thus we see that p is really close to zero. But this is clearly false since the largest power of two dividing $p(11) = (-2)(-4)(-6)(-8)(-10)$ is 2^8 . Thus we have a contradiction. Hence no such polynomial q exists, and thus we conclude that p is in fact partial credit. **CB: ACH,JVP**)

8. If $f(x) = x^{x^{x^x}}$, find the last two digits of $f(17) + f(18) + f(19) + f(20)$.

(ANS: 32. We are interested in finding $f(17) + f(18) + f(19) + f(20)$ modulo 100. By the Chinese Remainder Theorem it suffices to find the sum modulo 4 and 25. $A \equiv 0 \pmod{4}$ because 18 and 20 are raised to high powers and 19 and 17 are both raised to odd powers.

PUMaC 2008-9



Number Theory - A

$$18^4 \equiv 1 \pmod{25} \Rightarrow f(18) \equiv 1 \pmod{25}$$

$$f(20) \equiv 0 \pmod{25}$$

To compute $f(17)$ modulo 25, we need to compute $X = 17^{17^{17}}$ modulo 20. $X \equiv 1 \pmod{4}$ since $17 \equiv 1 \pmod{4}$. $X \equiv 2 \pmod{5}$ and therefore $X \equiv 17 \pmod{20}$. Since $3 * 17 \equiv 1 \pmod{25}$, $17^{17} \equiv 3^3 \pmod{25}$.

$$f(17) \equiv 2 \pmod{25}$$

To compute $f(19)$ modulo 25, we need to compute $Y = 19^{19^{19}}$ modulo 20.

$$19 \equiv -1 \pmod{20} \Rightarrow Y \equiv -1 \pmod{20}$$

$$f(19) \equiv 19^{-1} \equiv 4 \pmod{25}$$

$f(17) + f(18) + f(19) + f(20) \equiv 2 + 1 + 4 + 0 \equiv 7 \pmod{25}$.
 $f(17) + f(18) + f(19) + f(20) \equiv 0 \pmod{4}$ because 18 and 20 are raised to high powers and 19 and 17 are both raised to odd powers.
 $f(17) + f(18) + f(19) + f(20) \equiv 0 \pmod{4}$.

The last two digits of $f(17) + f(18) + f(19) + f(20)$ are 32. **CB: AL, IAF)**


9. What is the largest integer which cannot be expressed as $2008x + 2009y + 2010z$ for some positive integers x, y , and z ?

(ANS: 2016031. The largest number that cannot be a positive linear combination of 1004 and 1005 is $1004 * 1005 - 1004 - 1005 = 1007011$, so the smallest even number that cannot be written as the sum of 2008 and 2010 is 2007022. Then the largest number that we can't get is 2009031, as every larger number you can make it even by subtracting 2009 if necessary and write that as the sum of multiples of 2008 and 2010. **CB: AL12, IAF)**

10. Find the smallest positive integer n such that $32^n = 167x + 2$ for some integer x .

(ANS: 50. $32^{50} = 13^{500} = 13^{2+3*166} = 13^2 = 2$, as 167 is prime. Since 2 is a perfect square, and nothing else in the unit group, the answer is less than 83, and so it is 50. Alternately, $32 = 2^5$. $2^{83} \equiv 1 \pmod{167}$. So we need the smallest n such that $5n \equiv 1 \pmod{83}$. That gives $n = 50$. **CB: EK)**

PUMaC 2008-9



Number Theory - A

11. Find all sets of three primes p , q , and r such that $p + q = r$ and $(r - p)(q - p) - 27p$ is a perfect square.

(ANS: 2, 29, 31. It is easy to see that $q > p$ and that one of p and q is two, so $p = 2$. We then have $q(q - 2) - 54 = x^2$, so $(q - 1)^2 - x^2 = 55$. Thus, since the prime factorization of 55 is $5 * 11$, we have either $q - 1 = 8$ and $x = 3$, or $q - 1 = 28$ and $x = 27$. Only $q = 29$ has the solution be only primes. **CB:** AL, IAF)

12. Find the number of positive integer solutions of $(x^2 + 2)(y^2 + 3)(z^2 + 4) = 60xyz$.

(ANS: 8. We have that $xyz < 60$. We also have that, if a solution uses $z = 1$, then you can replace that with $z = 4$, and similarly, you can replace $y = 1$ with $y = 3$, and $x = 1$ with $x = 2$. Moreover, you have that $z \equiv \pm 1(5)$, as there is no way to force 5 to divide $x^2 + 2$ or $y^2 + 3$. Since any solution of the equation can be forced to have $x, y \geq 2$, we have to only check $z = 1, 4, 6, 9, 11, \text{ and } 14$. The only possible solution for $z = 14$ is $x = y = 2$ with the already stated equivalences, and one can check that that fails. If $z = 11$, we need 2 powers of 5 on the right side of the equation, which forces xyz to be too big. If $z = 9$, we have that $17|xy$ and so xyz is too big. If $z = 6$, we have that $(x^2 + 2)(y^2 + 3) = 9xy$. Thus $xy < 9$ and since $\nu_3(y) \geq \nu_3(y^2 + 3)$, we have $x = 4, 5$, and, since we can force y to be at least 2, we have $x = 4$ and $y = 2$, which doesn't work. Thus, we have $z = 4$ (or 1). The work here is similar and similarly tedious, and the only solution are the 8 that are equivalent to $x = y = z = 1$. **CB:** AL)

13. What is the smallest number n such that you can choose n distinct odd integers a_1, a_2, \dots, a_n , none of them 1, with $\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n} = 1$?

(ANS: 9. You can check that the answer is not less than 8, as $\frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{9} + \frac{1}{11} + \frac{1}{13} + \frac{1}{15} < 1$. You also can check that it is not 8, as the sum is, once you normalize denominators, the sum of odd numbers over an odd number. Thus, we need to show that 9 works. To see that 9 works, you can choose 3, 5, 7, 9, 11, 15, 35, 45, 231 as the a_i . **CB:** IAF)