




## Geometry B Solutions

- Find 100 times the area of a regular dodecagon inscribed in a unit circle. Round your answer to the nearest integer if necessary.



**Solution.** 300. The area is:

$$12 \times \frac{1}{2} \times 1^2 \times \sin 30^\circ = 3$$

- A triangle has sides of lengths 5, 6, 7. What is 60 times the square of the radius of the inscribed circle?

**Solution.** 160.


Use Heron's Formula, and then the area of the triangle is

$$\sqrt{9 \times (9 - 5) \times (9 - 6) \times (9 - 7)} = 6\sqrt{6}$$

Hence radius of incircle is

$$\frac{2 \times 6\sqrt{6}}{5 + 6 + 7} = \frac{2\sqrt{6}}{3}$$

- A rectangular piece of paper  $ABCD$  has sides of lengths  $AB = 1$ ,  $BC = 2$ . The rectangle is folded in half such that  $AD$  coincides with  $BC$  and  $EF$  is the folding line. Then fold the paper along a line  $BM$  such that the corner  $A$  falls on line  $EF$ . How large, in degrees, is  $\angle ABM$ ?





**Solution.** 30. Construct  $NG \perp BC$  at  $G$ . Triangles  $ABM$ ,  $NBM$ ,  $NBG$  are all congruent. Hence  $\angle ABM = 30^\circ$


4. Tetrahedron  $ABCD$  has sides of lengths, in increasing order, 7, 13, 18, 27, 36, 41. If  $AB = 41$ , then what is the length of  $CD$ ?

**Solution.** 13. By triangle inequality,  $AB + DB > 41$ , and  $AC + CB > 41$ . Hence, one of the pairs  $AD$ ,  $DB$  and  $\{AC, CB\}$  must be  $\{18, 27\}$ , the other pair contains 36. WLOG, let  $AC = 27$ ,  $CB = 18$ . Then  $DB \neq 36$ , otherwise,  $CD > 18$ . Hence  $AD = 36$ ,  $CD = 13$ .


5. A polygon is called concave if it has at least one angle strictly greater than  $180^\circ$ . What is the maximum number of symmetries that an 11-sided concave polygon can have?

**Solution.** 1. An 11-gon can have only axes passing through a vertex and an opposite side. If it had exactly 2 such axes, they would have to be perpendicular and the polygon would have an even number of sides. If it had exactly 3 axes, each would have to be at an angle of  $60^\circ$  with the next one, and the number of sides of the polygon would be a multiple of 3. Since our polygon has a concave angle, any number of axes of symmetry above 3 would imply at least 12 sides.

6. In the following diagram (not to scale),  $A, B, C, D$  are four consecutive vertices of an 18-sided regular polygon with center  $O$ . Let  $P$  be the midpoint of  $AC$  and  $Q$  be the midpoint of  $DO$ . Find  $\angle OPQ$  in degrees.



**Solution.** 30. Let  $R$  be the midpoint of  $AO$ . Connect  $RP$ ,  $RQ$ . Then  $RP = RO = OQ = RQ$ . Hence triangle  $PQR$  is isosceles, then some simple calculation yields  $\angle OPQ = \angle RPQ - \angle RPO = 50^\circ - 20^\circ = 30^\circ$ .



7. Lines  $l$  and  $m$  are perpendicular. Line  $l$  partitions a convex polygon into two parts of equal area, and partitions the projection of the polygon onto  $m$  into two line segments of length  $a$  and  $b$  respectively. Determine the maximum value of  $\lfloor \frac{1000a}{b} \rfloor$ . (The floor notation  $\lfloor x \rfloor$  denotes largest integer not exceeding  $x$ )

**Solution.** 2414. The greatest possible value of the ratio is  $(1 + \sqrt{2})$ . Let  $A$  and  $B$  be vertices of the convex polygon on different sides of  $l$  so that their distance from  $l$  is maximal on each side. Let  $K$  and  $L$  be the intersections of  $l$  with the sides of the polygon. Define the points  $K_1$  and  $L_1$  on extension of  $AK$ ,  $AL$  respectively such that  $K_1L_1$  is parallel to  $l$ . Since the polygon is convex, the part of the polygon on  $A$ 's side contains  $AKL$ , and the part of the polygon on  $B$ 's side is contained in  $K_1KLL_1$ . Therefore  $[K_1KLL_1] \geq [AKL]$ . Let  $M$ ,  $N$  be foot of perpendicular from  $A$ ,  $B$  to line  $l$  respectively, then

$$\begin{aligned} \frac{[AKL]}{[AK_1L_1]} &\leq \frac{1}{2} \implies \frac{AM}{AM + BN} \leq \frac{1}{\sqrt{2}} \\ &\implies \frac{AM}{BN} \leq \frac{1}{\sqrt{2} - 1} = 1 + \sqrt{2} \end{aligned}$$

Equality is obtained when the polygon is a triangle with  $l$  parallel to one side.

8. Consider the solid with 4 triangles and 4 regular hexagons as faces, where each triangle borders 3 hexagons, and all the sides are of length 1. Compute the *square* of the volume of the solid. Express your result in reduced fraction and concatenate the numerator with the denominator (e.g., if you think that the square is  $\frac{1734}{274}$ , then you would submit 1734274).

**Solution.** 52972. Extend the edges that are common to two hexagons. We obtain a regular tetrahedron of side length 3. Hence the volume of original solid is a regular tetrahedron of side length 3 minus volume of 4 regular tetrahedrons of side length 1. The volume is

$$\frac{1}{3} \times \frac{9\sqrt{3}}{4} \times \sqrt{6} \times \frac{27-4}{27} = \frac{23\sqrt{2}}{12}$$