## PUMaC 2011



## A MAN DOLLAR

## Algebra B

1. [3] If we define  $\otimes (a, b, c)$  by

$$\otimes(a,b,c) = \frac{\max(a,b,c) - \min(a,b,c)}{a+b+c - \min(a,b,c) - \max(a,b,c)},$$

compute  $\otimes(\otimes(7,1,3),\otimes(-3,-4,2),1)$ .

- 2. [3] If a and b are the roots of  $x^2 2x + 5$ , what is  $|a^8 + b^8|$ ?
- 3. [4] Let  $f(x) = x^3 7x^2 + 16x 10$ . As x ranges over all integers, find the sum of distinct prime values taken on by f(x).
- 4. [4] Let f be an invertible function defined on the complex numbers such that

$$z^{2} = f(z + f(iz + f(-z + f(-iz + f(z + \ldots)))))$$

for all complex numbers z. Suppose  $z_0 \neq 0$  satisfies  $f(z_0) = z_0$ . Find  $1/z_0$ . (Note: an invertible function is one that has an inverse).

5. [5] A polynomial p can be written as

$$p(x) = x^6 + 3x^5 - 3x^4 + ax^3 + bx^2 + cx + d.$$

Given that all roots of p(x) are equal to either m or n where m and n are integers, compute p(2).

- 6. [6] Shirley has a magical machine. If she inputs a positive even integer n, the machine will output n/2, but if she inputs a positive odd integer m, the machine will output m+3. The machine keeps going by automatically using its output as a new input, stopping immediately before it obtains a number already processed. Shirley wants to create the longest possible output sequence possible with initial input at most 100. What number should she input?
- 7. [7] A sequence of real numbers  $\{a_n\}_{n\geq 1}$  has the following property:

$$6a_n + 5a_{n-2} = 20 + 11a_{n-1}$$
 (for  $n \ge 3$ ).

The first two elements are  $a_1 = 0, a_2 = 1$ . Find the integer closest to  $a_{2011}$ .

8. [8] Let  $\alpha_1, \alpha_2, \ldots, \alpha_6$  be a fixed labeling of the complex roots of  $x^6 - 1$ . Find the number of permutations  $\{\alpha_{i_1}, \alpha_{i_2}, \ldots, \alpha_{i_6}\}$  of these roots such that if  $P(\alpha_1, \ldots, \alpha_6) = 0$ , then  $P(\alpha_{i_1}, \ldots, \alpha_{i_6}) = 0$ , where P is any polynomial with rational coefficients.