



## Algebra B Solutions

Written by Monica Marinescu

1. [3] Find the largest  $n$  such that the last nonzero digit of  $n!$  is 1.

Solution: For  $n > 2$  if we express  $n! = 2^i 5^j m$  where,  $i, j, m \in \mathbb{N}$  such that  $m$  is not divisible by 2 or 5, we can easily observe that  $i > j$ . Thus, we can write  $n! = 2^{i-j} 10^j m$  which means that the last non-zero digit of  $n!$  is always going to be an even number. Thus, the only possible values for  $n$  are 0 and 1, which both give the last non-zero digit equal to 1. Because 1 is greater, the final answer is 1.

Answer: 1

2. [3] Define a sequence  $a_n$  such that  $a_n = a_{n-1} - a_{n-2}$ . Let  $a_1 = 6$  and  $a_2 = 5$ . Find  $\sum_{n=1}^{1000} a_n$ .

Solution:  $a_1 = 6, a_2 = 5, a_3 = -1, a_4 = -6, a_5 = -5, a_6 = 1, a_7 = 6, a_8 = 5$ , so the sequence starts to repeat itself. The sum of the first 6 terms is 0, so the sum of the first 996 terms is 0; thus, the sum of the first 1000 terms is  $6 + 5 + (-1) + (-6) = 4$ .

Answer: 4

3. [4] Evaluate  $\sqrt[3]{26 + 15\sqrt{3}} + \sqrt[3]{26 - 15\sqrt{3}}$ .

Solution: Let  $x = \sqrt[3]{26 + 15\sqrt{3}}, y = \sqrt[3]{26 - 15\sqrt{3}}$ . We are looking for  $z = x + y$ . We can do this by cubing  $x + y$ .

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x + y)^3 = 26 + 15\sqrt{3} + 26 - 15\sqrt{3} + 3\sqrt[3]{26^2 - 3 * 15^2}(x + y)$$

$$(x + y)^3 = 52 + 3\sqrt[3]{1}(x + y)$$

$$z^3 = 52 + 3z$$

$$z^3 - 3z - 52 = 0$$

$$(z - 4)(z^2 + 4z + 13) = 0$$

Since  $z^2 + 4z + 13$  has no real root,  $z$  has to equal 4.

Answer: 4

4. [4] If  $x, y$ , and  $z$  are real numbers with  $\frac{x-y}{z} + \frac{y-z}{x} + \frac{z-x}{y} = 36$ , find

$$2012 + \frac{x-y}{z} \cdot \frac{y-z}{x} \cdot \frac{z-x}{y}$$



Solution:

$$\begin{aligned}
 36 &= \frac{(x-y)xy + (y-z)yz + (z-x)xz}{xyz} \\
 &= \frac{(x-y)xy + y^2z - yz^2 + xz^2 - x^2z}{xyz} \\
 &= \frac{(x-y)xy - (x+y)(x-y)z + z^2(x-y)}{xyz} \\
 &= \frac{(x-y)(xy - xz - yz + z^2)}{xyz} \\
 &= \frac{(x-y)(y-z)(x-z)}{xyz} \\
 36 &= -\frac{(x-y)(y-z)(z-x)}{xyz} \\
 2012 + \frac{(x-y)(y-z)(z-x)}{xyz} &= 2012 - 36 = 1976
 \end{aligned}$$

Answer: 1976

5. [5] Considering all numbers of the form  $n = \lfloor \frac{k^3}{2012} \rfloor$ , where  $\lfloor x \rfloor$  denotes the greatest integer less than or equal to  $x$ , and  $k$  ranges from 1 to 2012, how many of these  $n$ 's are distinct?

Solution:

$$\begin{aligned}
 \frac{(k+1)^3}{2012} - \frac{k^3}{2012} &= \frac{3k^2 + 3k + 1}{2012} \\
 \frac{3k^2 + 3k + 1}{2012} &\geq 1 \iff 3k^2 + 3k + 1 \geq 2012 \\
 \iff 3k^2 + 3k &\geq 2011 \iff 3k^2 + 3k \geq 2010 \\
 \iff k^2 + k &\geq 670 \iff k \geq 26
 \end{aligned}$$

Thus, for  $k$  of at least 26, the difference between two consecutive fractions is at least 1, so the difference between their integer parts is also at least 1, so the numbers are different; in conclusion, for  $k$  between 26 and 2012, there are 1987 different numbers.

For  $k$  less than 26, the difference between two consecutive fractions is less than 1, so the integer parts of two consecutive fractions is at most 1. For  $k = 1$ ,  $\lfloor \frac{k^3}{2012} \rfloor = 0$ , and for  $k = 25$ ,  $\lfloor \frac{k^3}{2012} \rfloor = 7$ , so for  $k$  less than 26, there are 8 different numbers.

In the end, we get that for  $k$  ranging from 1 to 2012, there are 1995 different numbers in the sequence.

Answer: 1995

6. [6] Compute

$$\sum_{n=1}^{\infty} \frac{n+1}{n^2(n+2)^2}$$



Your answer in simplest form can be written as  $a/b$ , where  $a, b$  are relatively-prime positive integers. Find  $a + b$ .

Solution: Let  $K$  be the result.

$$\begin{aligned} K &= \sum_{n=1}^{\infty} \frac{n+1}{n^2(n+2)^2} \\ 4K + \sum_{n=1}^{\infty} \frac{1}{(n+2)^2} &= \sum_{n=1}^{\infty} \frac{n^2 + 4n + 4}{n^2(n+2)^2} \\ 4K - 1 - \frac{1}{4} + \sum_{n=1}^{\infty} \frac{1}{n^2} &= \sum_{n=1}^{\infty} \frac{(n+2)^2}{n^2(n+2)^2} \\ 4K - \frac{5}{4} + \sum_{n=1}^{\infty} \frac{1}{n^2} &= \sum_{n=1}^{\infty} \frac{1}{n^2} \\ 4K - \frac{5}{4} &= 0 \\ K &= \frac{5}{16} \end{aligned}$$

So  $a + b = 5 + 16 = 21$ .

Answer: 21

7. [7] Let  $f$  be a polynomial of degree 3 with integer coefficients such that  $f(0) = 3$  and  $f(1) = 11$ . If  $f$  has exactly 2 integer roots, how many such polynomials  $f$  exist?

Solution: The answer is 0, but the argument is more general: if  $f(0)$  and  $f(1)$  are odd, then we claim that  $f$  can't have any integer roots:

Suppose  $a$  is an integer solution. Then  $f(x) = (x-a)g(x)$ , and  $g(x)$  also has integer coefficients. So  $f(0) = -ag(0)$  and  $f(1) = (1-a)g(1)$ , where  $g(0)$  and  $g(1)$  are also integers. Since either  $a$  or  $1-a$  is even,  $f(0)$  and  $f(1)$  can't be both odd, as in the hypothesis, so  $f$  has no integer roots.

Answer: 0

8. [8] Let  $a_n$  be a sequence such that  $a_1 = 1$  and  $a_{n+1} = \lfloor a_n + \sqrt{a_n} + \frac{1}{2} \rfloor$ . What are the last four digits of  $a_{2012}$ ?

Solution: Computing some particular cases suggests that the function  $f$  is defined by the following:

$$a_n = 1 + \lfloor \frac{n}{2} \rfloor \lfloor \frac{n+1}{2} \rfloor,$$

for all natural numbers  $n$ .

We will show this hypothesis is true by induction. We assume it's true for  $n$  and prove it for  $n+1$ .



If  $n = 2m$ , then  $a_n = 1 + m^2$ . Thus,

$$\begin{aligned} a_{n+1} &= \lfloor a_n + \sqrt{a_n} + \frac{1}{2} \rfloor = \lfloor m^2 + 1 + \sqrt{m^2 + 1} + \frac{1}{2} \rfloor \\ &= m^2 + 1 + m = 1 + m(m+1) = 1 + \lfloor \frac{n+1}{2} \rfloor \lfloor \frac{n+2}{2} \rfloor \end{aligned}$$

If  $n = 2m + 1$ , then  $a_n = 1 + m(m+1)$ , thus

$$\begin{aligned} a_{n+1} &= \lfloor 1 + m(m+1) + \sqrt{1 + m(m+1)} + \frac{1}{2} \rfloor = 1 + m(m+1) + m + 1 \\ &= 1 + (m+1)^2 = 1 + \lfloor \frac{n+1}{2} \rfloor \lfloor \frac{n+2}{2} \rfloor \end{aligned}$$

The induction is complete, so that is the solution to the equation. In conclusion,  $a_{2012} = 1 + \lfloor \frac{2012}{2} \rfloor \lfloor \frac{2013}{2} \rfloor = 1012037$ , so the answer is 2037.

Answer: 2037