训励新林等席 State Mathematics Algebra II Contest May 3, 2001 Solutions

mutall m # "

Institute \$ 75

Y.

Y.

N.

Y.

N.

Y.

Ro

to the the B. Ph

to the the B Ph

(L+1)(W+1) = 84 and (L-1)(W-1) = 48 so we get LW+L+W+1 = 841. (c) LW - L - W + 1 = 48Subtracting these last two equations yields 2L + 2W = 36, so we could go on and solve

Institute # # 3

而时间的称林塔梯

Institute \$ * 3

moutule # # 3 PS

to the the the

to the the B Ph

Institute mark 'S

Artitute the the 'S PR

Institute m # "

stitute # # # K for L and W, however, we are looking for the perimeter, so we are done. The terms of the sequence, all in terms of w and x are 2. (e)

 $\dots, w, x, w + x, w + 2x, 2w + 3x, 3w + 5x, \dots$ Thus 2w + 3x = 0 and 3w + 5x = 1. Solving this system of equations gives w = -3, x = 2, so the terms are $\dots, -3, 2, -1, 1, 0, 1, \dots$, 加加新林塔梯 making the quantity 2(w + x + y + z) = 2(-3 + 2 - 1 + 1) = -2. 恢秋

加斯林塔张 4. % W (b) The only case not to be considered is if both are tails, and this occurs with a \$50 probability of (0.75)(0.40) = 0.30. Thus there will be at least one head 70% of the time.

5. (a) If three of the zeros are 2, 1+i, 3-i, and the coefficients are real, then the conjugates of the imaginary roots must also be zeros. Thus the minimum degree of the polynomial will be five. astitute that the 'S PR

mstitute # 6. With the three points (0,-1), (1,4) and (2,13), we could find all of the (d) coefficients of the quadratic, however, since we are looking for the sum of the coefficients, we can plug (1,4) into $y = ax^2 + bx + c$ and get $4 = a \cdot 1^2 + b \cdot 1 + c$.

 $8^{6x^2+4x} = 4^{9x^2-9x+6} \Rightarrow 2^{3(6x^2+4x)} = 2^{2(9x^2-9x+6)}$. Now with the bases the same withthe start " S PR $\Rightarrow 2$ $= 2^{-(xx^2-yx+0)}$. Now with the bases the same we know that the exponents must be the same, so $3(6x^2+4x) = 2(9x^2-9x+6)$. This simplifies to 30x - 12 of $x = 2^{1/3}$ simplifies to 30x = 12 of $x = \frac{2}{5}$.

8. (b) Looking at the amount of grape juice, we get $d \cdot \frac{d}{100} + x = (d + x)\frac{3d}{100}$, where x is the amount of pure grape juice to be added. This simplifies to ansitute ## # 13 PR $d^{2} + 100x = 3d^{2} + 3dx$. Solving for x we get $x = \frac{2d^{2}}{100 - 3d}$.

9. The following sequence of graphs gives a progression that leads directly to 面对加根教林後隊 (c) no. 28 8 4 18 million mailule # # % K matitute ## # 'E K Withite # # 12 1% the answer. n ****

to the bit of the



multitute # # 18. time # # B One extreme for the shortest altitude will occur when the two R (d) shortest altitudes are both 12, in an isosceles triangle. The other extreme occurs in a figure similar to the one below. From similar triangles в z Ν 12, Multilite # # 3 motilite # # 3 PR multille # # 'E K w Y. h х 11x 2 66 motilite # # 13 PR multitute ## # '\$ PR O to H & R multitute ## # '3 PR $\triangle BMA \simeq \triangle COA \text{ that } \frac{12}{x} = \frac{66}{AO} \Rightarrow AO = \frac{11x}{2}. \text{ Also from similar}$ triangles $\triangle BMC \simeq \triangle ANC$, we know that $\frac{12}{x+u} = \frac{h}{w} \Rightarrow h = \frac{12w}{x+u}$. But we 面对机机新林塔像 know that u > w and $\frac{11x}{2} < u$, so $h = \frac{x+u}{x+u} < \frac{w}{x+u} < \frac{x+u}{x+u}$. Simplifying N. this we have $h < \frac{12u}{\frac{13u}{2}} = \frac{132u}{13u} = 10\frac{2}{13}$. Thus $12 \le x < 10\frac{2}{3}$, and the only mutule ## # '& R distinct integer length for the shortest altitude is 11. Let M and N be the two numbers. The problem states that M must o digit number, but does not indicate the N the number 19. (c) be a two digit number, but does not indicate the N, the number formed when the digits are reversed, also must be a two digit number. So, let M = 10a + b, where $0 < a \le 9$ and $0 \le b \le 9$. Then N = 10b + a, and the mutute # # B 加度频频发展 difference M - N = (10a + b) - (10b + a) = 9(a - b). Now for *M* - *N* to be a perfect cube, a - b must be a multiple of 3, so we have, for M, 96, 85, 74, 63, 52, 41, and 30. 20. We find that side AC and AB both have length 5, so the angle-(e) bisector is also the median, passing through A(1,-2) and the midpoint mythute ## # '& PL 面的地格林塔塔 (3/2,3/2), of side BC. This makes the slope 7 and the equation y = 7(x-1) - 2 = 7x - 9. 21.titute To find the inverse, exchange the x and the y and solve. (b) $x = \frac{3y-5}{2y+1} \Rightarrow 2xy + x = 3y-5 \Rightarrow (2x-3)y = -x-5$, so $y = \frac{x+5}{-2x+3}$, 而此此他新林塔像 mstitute ## # 18 而如此他教祥学家 matinue ## # '\$ 1% making the product $5 \cdot (-2) \cdot 3 = -30$ Y. Institute # 3

multille m # "

Y.

Y.

to the the B

Ro

to the We the Ph

to the W- 1/3 Ph

to the We B

multille m # *

Institute m # "

multine m # 3

multinu m # "

multille m # 3

to the We By Pho

to the W- 1/2 Ph

N.

multille # # 3 PR J. (a) H 3 IF If you expand the expression $(\sqrt[3]{4} + \sqrt[3]{2} - 2)(a\sqrt[3]{4} + b\sqrt[3]{2} + c) = 20$, you get $2^{\frac{2}{3}}(c+b-2a)+2^{\frac{1}{3}}(c+2a-2b)+2(a+b-c)=20$. Since we know that a, b, and c are integers, the first two terms must be zero, and the final term, 2(a+b-c) must equal 20, so a+b-c=10. Note: You can Institute # # 'S PE Withthe Start H 13 PR solve for a, b, and c, getting (a,b,c) = (6,8,4), but there is no need to since the problem asks for a+b-c. 31. (e) If we write the fractions in order as $\binom{1}{2} + \binom{1}{3} + \binom{2}{3} + \binom{1}{4} + \binom{2}{4} + \binom{3}{4} + \dots + \binom{1}{100} + \binom{2}{100} + \dots + \binom{99}{100}$, we Institute # # 3 PS see that this sum is $\binom{1}{2} + \binom{3}{3} + \binom{6}{4} + \binom{10}{5} + \binom{15}{6} + \dots + \binom{\frac{10099}{2}}{100}$, using the formula for the sum of n-consecutive integers $1+2+3+\dots+n=\frac{n(n+1)}{2}$. If you now factor one-half from each term you 而如此他就林塔张 get $\frac{1}{2}(1+2+3+4+\dots+99) = \frac{1}{2}\left(\frac{99\cdot100}{2}\right) = 2475$. 32.^{titut} Since we know that the area is $a^2 + b^2 - c^2$, we know that *c* cannot (c) be the hypotenuse, it must be a or b. We know from the question that a is 而此此他新祥後 the hypotenuse. Now we know that $b^2 + c^2 = a^2$ and that the area is $\frac{1}{2}bc$, so $\frac{1}{2}bc = a^2 + b^2 - c^2 \Rightarrow \frac{1}{2}bc = 2b^2$ since $a^2 - c^2 = b^2$, Thus $\frac{c}{b} = 4$. $2 - \frac{2}{5}t = 1 - \frac{1}{7}t \Leftrightarrow 1 = \frac{2}{5}t - \frac{1}{7}t = \frac{9}{35}t$. So $t = \frac{35}{9} = 3.\overline{8}$ hours. 34. (b) Since the Let t be time, so we are told that $2(1-\frac{1}{5}t) = 1(1-\frac{1}{7}t)$. Thus 面前加速新林塔梯 Since the remainder is a constant, P(x) must be quadratic. Look first at the simple quadratic $P(x) = x^2 + bx + c$. Since x + 1 yields the remainder of -5 when divided into P(x), we know that 面射机机新林塔张

minitule # # *

mistille # **

Institute \$ ****

to the the the the

multille m # "

Y.

Y.

N.

而此此他新林塔像

古田林後代

** 标林 塔

Y.

Y.

multilite # # *

militule # ** *

P(-1) =1 -b +c = -5. We also know that P(5) = 25+5b+c = 7. Solving for b and c we get b = -2 and c = 8. Now the product $x^2 - 4x - 5$ divides into $P(x) = x^2 - 2x - 8$ one time with a remainder of 2x - 3. Note: If you use a more complicated quadratic for P(x), so $P(x) = ax^2 + bx + c$, you find that any *a* will work, and with b = 2 - 4a and c = -3 - 5a, the product $x^2 - 4x - 5$ divides into P(x) a times with the same remainder.

1. 长 长

~ 资本

35. (a) Complete the squares to find the center of the circle. $x^2-6x+9+y^2-4y+4=-11+9+4=2$, so $(x-3)^2+(y-2)^2=2$, putting the center of 2.2 Matilute # # 35. Y. putting the center at (3,2). This point is $\sqrt{(3-0)^2 + (2-0)^2} = \sqrt{13}$ from the origin. 面动机机称林塔梯 36. (a) $\log_2(a^3b) = x \Leftrightarrow 3\log_2 a + \log_2 b = x$. $\log_2\left(\frac{3a}{b}\right) = y \Leftrightarrow \log_2 3 + \log_2 a - \log_2 b = y$. Adding these last two expressions give us $4\log_2 a + \log_2 3 = x + y$. Solving for $\log_2 a$ yields 面动机机都林塔梯 面前加速素林等除 $\log_{2} = \frac{x+y}{4} - \frac{1}{4}\log_{2} 3 = \frac{x+y}{4} - \log_{2} \sqrt[4]{3}.$ 37. (c) $2x + 3y = A \\ x + 2y = B \Leftrightarrow 2x + 3y = A \\ 2x + 4y = 2B \Rightarrow y = 2B - A.$ But Y. $x = B - 2y \Longrightarrow x = B - 2(2B - A) = 2A - 3B.$ 而如此他就林塔张 (d) When you expand $(a+b)^8$ you get a nine terms, each with some Y. 38. coefficient and a and b raised to powers from 0 to 8. Plugging 1 in for both a and b would then give you the sum of the coefficients, to go ahead and plug 1 in at the beginning, getting $(1+1)^8 = 2^8 = 256$. multille # # 3 PS Y. (d) To find the inverse, exchange the x and y and solve for y. Thus $x = \frac{y+1}{y-1} \Leftrightarrow xy - x = y+1 \Leftrightarrow xy - y = x+1 \Leftrightarrow y = \frac{x+1}{y-1}$. From this we see that the function is its own inverse. If we had graphed it, and noticed the symmetry about the line y = x, we could have drawn the same conclusion. 而如此他教林塔梯 astitute # # B PK Thus $f^{-1}\left(\frac{1}{x}\right) = f\left(\frac{1}{x}\right) = \frac{\frac{1}{x}+1}{\frac{1}{x}-1} = \frac{x+1}{x-1}$. Setting this equal to 3, we have Y. $\frac{x+1}{x-1} = 3 \Leftrightarrow x+1 = 3(x-1) \Rightarrow 2x = 4, \text{ so } x = \frac{1}{2}.$ $\frac{k-3}{0-4} = \frac{3-0}{4-p} \Rightarrow (k-3)(p-4) = 12$. The positive integer factors of 12 are Let the y-intercept by (0,k) and the x-intercept (p,0). Then N. $p-4=1,2,3,4,6,12 \Rightarrow p=5,6,7,8,10,16$ 1, 2, 3, 4, 6, and 12, so $k-3=12,6,4,3,2,1 \Rightarrow k=15,9,7,6,5,4$. Of Inditute # # '& R multure the tot 'S PR Withite the the 's PR matitute ## # 18 Y. these, only p = 5,7 are prime. rs مراجع stitute # to the the B to the With the Pho to the the 'S Ph 小 妆 妆 客 粥 to the the 1/2 1/2 to the We B Ro

multille m # "

Institute # ** *

Institute m # "

Institute m # *

mutall m # 3

mating # **